

$S_2S_3S_4 = 107^\circ$ and $S_2S_3S_4/S_3S_4S_5 = 106^\circ$. The slight differences between these values and those reported earlier² for the orthorhombic structure, are within the experimental errors.

The triclinic and orthorhombic crystals both have a layer structure, and show a corresponding perfect cleavage along the *c* plane. The thickness of the layers are, $d_{001} = 10.78 \text{ \AA}$ and half the orthorhombic *c* axis, *viz.*, 10.89 \AA , respectively. Within the probable errors of the structure determinations, the atomic arrangement *within* the layers is the same in the two crystals. The orthorhombic space group requires a mirror plane of symmetry to pass through the barium ion and the middle sulphur atom of the pentathionate chain. Although not crystallographically required, a mirror plane of symmetry is actually present in the layers of the triclinic crystals, and is depicted through broken lines in Fig. 1. The plane is normal to, and passes through, the *b* axis at $z = 0$, $y = \frac{1}{2}$ and $\frac{3}{2}$, as in the orthorhombic crystals, and through the same atoms. The orthorhombic and triclinic modifications differ only in the arrangement of the layers relative to each other.

A detailed account of the structure will be published later.

The work has been aided by grants from *Norges Almenvitenskapelige Forskningsråd* and from *Norges Tekniske Högskoles Fond*.

1. Foss, O. *Acta Chem. Scand.* **7** (1953) 697.
2. Foss, O. and Zachariassen, H. *Acta Chem. Scand.* **8** (1954) 473.

Received June 17, 1955.

Plant Growth Regulators I.

1- and 2-Naphthylmethylarsonic Acids

SVEN-OLOV LAWESSON

Department of Organic Chemistry, Chemical Institute, University of Uppsala, Uppsala, Sweden

In order that a substance may exhibit auxin activity it must, among other things, have an unsaturated ring system and an acidic side chain. Hitherto mostly synthetic plant hormones with a carboxyl group have been investigated. Following a suggestion by Professor A. Fredga the author has started an investigation on

Acta Chem. Scand. **9** (1955) No. 6

aromatic arsonic acids which, if physiologically active, will be of interest stereochemically and in interpreting the growth regulating mechanism.

The 1-naphthylmethylarsonic acid (I) and the 2-naphthylmethylarsonic acid (II) were prepared according to the general procedure outlined by Quick and Adams¹. The yields were very low but can certainly be increased. All analyses for arsenic were performed by the method described by Ramberg and Sjöström² and the titrimetric determinations in accordance with a method by King and Rutterford³.

The biological activity is being investigated by Dr. Börje Åberg, who has kindly reported some preliminary results. Both acids show in different tests a conspicuous anti-auxin effect⁴.

Experimental. 1-Naphthylmethylarsonic acid was prepared from 1-naphthylmethylchloride following the method given for benzylarsonic acid by Quick and Adams¹. Colourless needles. M. p. $142-144^\circ$ (decomposition). Calc. for $C_{11}H_{11}O_3As$ (266.1): C 49.6; H 4.17; As 28.2; equiv. wt. 133.1. Found: C 49.9; H 4.13; As 28.1; equiv. wt. 132.9.

2-Naphthylmethylarsonic acid was prepared in the same way from 2-naphthylmethyl bromide. Colourless plates. M. p. $159-161^\circ$ (decomposition). Found: C 50.0; H 3.99; As 28.0; equiv. wt. 133.8. Calc. for $C_{11}H_{11}O_3As$ (266.1): C 49.6; H 4.17; As 28.2; equiv. wt. 133.1.

1. Quick, A. J. and Adams, R. *J. Am. Chem. Soc.* **44** (1922) 805.
2. Ramberg, L. and Sjöström, G. *Svensk Kem. Tidskr.* **29** (1917) 73.
3. King, H. and Rutterford, G. V. *J. Chem. Soc.* **132** (1930) 2138.
4. Åberg, B. *Private communication*.

Received July 5, 1955.

Halogenated Guaiacoxylalkylcarboxylic Acids of Plant Physiological Interest

MAGNUS MATELL

Department of Organic Chemistry, Royal Agricultural College, Uppsala, Sweden

If the ether linkage in aryloxyalkylcarboxylic acids (I) is replaced by S, NH or CH_2 the plant growth-regulating activity is decreased^{1,2}. This type of