Short Communications

Acid Hydrolysis of Some α,β -Unsaturated Aliphatic Esters

ERRO A. HALONEN

Institute of Chemistry, University of Helsinki, Finland

The ethyl esters of propiolic, tetrolic and *tert.* butylpropiolic acid and the diethyl ester of acetylenedicarboxylic acid were prepared previously¹. In the present study, the hydrolyses were carried out in ampoules of Pyrex glass and of about 10.5 ml capacity. The sample volume was 10 ml and the concentrations of sulphuric acid and ester were both 0.02 N in the initial mixture. The reaction was arrested in ice and the acid was titrated with barium hydroxide using phenolphthalein as indicator.

Addition of water to the triple bond of the esters was not observed. If it occurred, ethyl propiolate would have given, *via* acetoacetic acid, acetaldehyde, and, correspondingly, ethyl tetrolate and ethyl *tert.* butylpropiolate would have given ketones. The acid hydrolysis of diethyl acetylenedicarboxylate cannot be studied by using the above mentioned method as the acidic ester resulting from the hydrolysis is decarboxylated at a faster rate than the diethyl ester is hydrolyzed. Also the propiolic acid formed in the hydrolysis of

ethyl propiolate is decarboxylated under these conditions² and, as a result, a decrease in the rate constant of hydrolysis occurs when the reaction proceeds. Especially at high temperatures, this becomes evident, as the activation energy of decarboxylation is 13 600 cal greater than the activation energy of hydrolysis. Mathematically, the decrease has been shown to agree, quantitatively, with the decarboxylation. The rate constants and the calculated activation energies and frequency factors are given in Table 1.

The ratios of the rates of hydrolysis correspond approximately to those previously obtained in alkaline hydrolysis of these esters¹.

The difference in the rates for ethyl propiolate and ethyl tetrolate is greater than the difference in the rates for ethyl tetrolate and ethyl *tert.* butylpropiolate. When compounds with triple bonds are compared to the corresponding saturated ones³ and to those with double bonds⁴, it will be noted, that the acetylenic compounds will hydrolyze about 40 times slower than the saturated ones but only slightly slower than the ethylenic compounds. The activation energies of the acetylenic and saturated compounds are of the same magnitude and approximately 3 000 cal smaller than the activation energies of the ethylenic compounds. This indicates that the conjugation between the unsaturated bond and the carboxyl carbonyl group has more effect

Table 1. Acid hydrolysis of esters in water and in acetone-water mixtures.

Ester	Solvent composition wt % H ₂ O	10 ⁴ k 1 mole ⁻¹ sec. ⁻¹			E cal	log A
		40.0°	60.0°	80.0°		
HC≡C·CO ₂ Et	100	1.16	5.60	22.4	16 221	4.39
→	50	0.437	2.03	7.74	15 680	3.59
H ₃ C·C≡C·CO ₂ Et	100	0.713	3.83	16.4	17 220	4.88
→	50	0.300	1.31	5.62	16 113	3.73
(CH ₃) ₃ C·C≡C·CO ₂ Et	50	0.276	1.14	4.53	15 860	3.46

on the rate of the hydrolysis than has the inductive effect and, on the basis of the reaction mechanism, this can also be expected. As compared with the ethylenic esters the decrease in the rate is due to the inductive effect of the triple bond. As a result of this either the pre-equilibrium moves from the side of the ionic conjugate acid of ester to the side of the ester or the alkoxyl linkage, after the addition of water, is less readily breakable.

The rates of hydrolysis in 60 % aqueous ethanol at 60° C for the ethyl ester of acetic, monochloroacetic, dichloroacetic and trichloroacetic acids are 1, 0.91, 0.40 and 0.11, respectively, if the rate for ethyl acetate is taken as a unity⁵. The fact that the rates of hydrolysis for the ethyl ester of propionic and acrylic acids differ only slightly from each other gives support to the assumption that, in the case of chloroacetic acids, steric factors are of importance.

1. Halonen, E. A. *Acta Chem. Scand.* **9** (1955) 1492.
2. Halonen, E. A. *Ann. Acad. Sci. Fennicae A. II.* **55** (1954).
3. Tommila, E. and Hella, A. *Ann. Acad. Sci. Fennicae A. II.* **53** (1954).
4. Pekkarinen, L. *Ann. Acad. Sci. Fennicae A. II.* **55** (1955).
5. Timm, E. W. and Hinshelwood, C. N. *J. Chem. Soc.* **1938** 862; Tommila, E., Hietala, S. and Nyrkiö, J. *Suomen Kemistilehti B.* **28** (1955) 143; Ingold, C. K. *Structure and Mechanism in Organic Chemistry*, Cornell University Press, New York 1953, p. 777.

Received September 20, 1956.

Paper Chromatographic Separation of Thiamine Phosphates

KARL-HEINZ KIESSLING

Institute of Zoophysiology, Uppsala, Sweden

Several papers have been published about paper chromatographic separation of thiamine phosphates¹⁻⁵. In all solvents described thiamine triphosphate hardly moves at all, and its R_F value coincides in most cases with that of nucleotide di- and triphosphates.

In this paper a paper chromatographic solvent is described in which thiamine tri-

phosphate moves easily, and in which it separates well from other thiamine compounds and from nucleotide phosphates. Also thiamine phosphates with more than three phosphate groups are separated. From a mixture of thiamine phosphates prepared according to Viscontini, Bonetti, and Karrer⁶, nine thiamine compounds can be isolated.

The solvent consists of 100 ml isobutyric acid, 60 ml N ammonia, and 1.6 ml 0.1 M ethylene diamine tetraacetic acid. It has been used by Krebs and Hems for separating nucleotides⁷ and is a modification of a solvent described by Zetterström and Ljunggren⁸. Filter papers Whatman No. 1 are used after washing in aqueous ethylene diamine tetraacetic acid according to the method described by Eggleston and Hems⁹.

The di- and triphosphates of inosine, guanosine, uridine, and cytidine all have lower R_F values than adenosine diphosphate¹⁰. In this investigation thiamine triphosphate was shown to move faster than adenosine diphosphate. It can thus be easily separated from the nucleotides mentioned.

Fig. 1 and 2 show the separation of different amounts of thiamine phosphates prepared according to Viscontini *et al.*⁶ One mg of the mixture dissolved in water is linearly spread on the filter paper in Fig. 1. The separation is very distinct. Ten lines can be seen with the naked eye, but only nine are registered on the photographic plate. When the amount of the thiamine phosphate mixture is increased to 3 mg (Fig. 2) the separation is still distinct except between thiamine mono- and diphosphate (lines 3 and 4). Nine lines can be detected.

The lines are cut out, the compounds run off with water, then added as spots on a new filter paper, and developed in the same solvent. The compounds appear as distinct spots with about the same R_F values as the lines on the original paper chromatogram (Fig. 3).

Total and hydrolyzable phosphorus per molecule of thiamine for the compounds 1-7 are seen in Table 1. Thus compound 1 and 2 were identified as different forms of thiamine*. The ratio between total and hydrolyzable phosphorus and thiamine suggests that the compounds 3-7 are thiamine mono-, di-, tri-, tetra- and pentaphosphate, respectively. Compounds

* These two forms have been separated earlier by Gaudiano, Spadoni and Tecce¹¹.