are also self-quenching. The colorless crystals did not fluoresce.

The spectrum of the white crystals (CHCl_3) in the infra red region showed prominent absorption peak at $2\,940\text{ cm}^{-1}$ with a shoulder at $2\,870\text{ cm}^{-1}$. Less prominent peaks are at $1\,725$, $1\,605$, $1\,467$, $1\,380$, $1\,060$, $1\,035$, 985 and 842 cm^{-1} . The purified green material exhibited the same prominent peak at $2\,920\text{ cm}^{-1}$ and the shoulder at $2\,860\text{ cm}^{-1}$, and in addition, the absorption at $1\,730\text{ cm}^{-1}$ was very much intensified. There was a great deal of generalized absorption from 900 — $1\,500\text{ cm}^{-1}$, with two distinct peaks at $1\,465\text{ cm}^{-1}$ and $1\,385\text{ cm}^{-1}$. The spectra are not overly informative, but do suggest the possible presence of aldehyde or COOH groups in the green compound, while the white crystals could be a saturated lipid having comparatively few functional groups.

The question whether the green compound is a natural constituent with a role in the physiology of the yeast cell cannot be answered unequivocally. It can be stated that the green compound was consistently obtained from each of five different air-dried brewers yeast preparations in this laboratory. This procedure, however, did not yield a green product from bakers yeast. It is pertinent, however, that Keilin³ reported a green fraction in initial attempts at the isolation of cytochrome *c* from brewers yeast in 1931. Efforts to obtain the present green compound by a suitable modification of Keilin's procedure have not proved successful.

There is some evidence which suggests that some of the efficiency of extraction of the green compound is very closely related to the state of the cells. Thus it has not been possible to obtain the green compound by extraction of fresh yeast cells, nor of cells which have been autolyzed even for short periods. It has also not been possible to obtain the green compound from freeze-dried preparations. Suitable experiments indicate that the green color is not due to an artifact of the reagents used.

The actual chemical structure of the green compound and its role in biological systems awaits clarification. It would be particularly interesting to know if present compounds have a relation to any of the physiologically important green compounds known.

Acknowledgment. The authors wish to express appreciation to Professor Theorell for his active interest in this work, and W. J. Rutter wishes to express his thanks for the gracious hospitality extended by Professor Theorell and other members of the Biochemical Department.

1. Lemberg, R. *Hematin Compounds and Bile Pigments*, Interscience Publishers, New York 1949.
2. Rabinowitch, E. I. *Photosynthesis*, Interscience Publishers, New York 1945—1951.
3. Keilin, D. *Proc. Roy. Soc. London B* **106** (1930) 418.

Received December 19, 1957.

Examination of the Behaviour of Liquid Mixtures in the Critical Region

HANS LUDVIG LORENTZEN and
BJØRN BERGSSNOV HANSEN

*Universitetets Kjemiske Institutt,
Blindern-Oslo, Norway*

In order to investigate the shape of the co-existence curves close to the critical point, the two systems of partly miscible liquids, triethylamine-water (at the lower critical point) and succinonitrile-ethanol have been examined. The results are shown in Fig. 1, where the co-existence curves are seen to follow the "cubic rule"^{1,2}

$$\Delta c = k \cdot \sqrt[3]{\Delta T} \quad (1)$$

Δc being the composition of the individual phases measured from the critical, and ΔT the temperature distance from the critical.

It will be observed from Fig. 1 that in the case of triethylamine-water the plotted results follow the straight line as close as 1 mdeg to the critical. However, uncertainties regarding the fixation of the critical temperature involve uncertainty as regards the plots from the critical temperature to about $\Delta T = 2\text{ mdeg}$. Nevertheless, the visually obtained information about the initiation of the two phases when passing the critical temperatures

Fig. 1. Top of co-existence curves. C_c is the critical concentration. T_c is the critical temperature. Curves are obtained for the critical systems succinonitrile-ethanol and triethylamine-water. The co-existence curve for the latter is obtained through series of experiments, each series giving four to eight points on the curve.

allows the assumption that the co-existence curve is regular to ΔT , this being only a fraction of a millidegree.

It should be noted that the critical temperature was not found to be constant. In the triethylamine-water system the critical temperatures seemed to have some connection with the initial concentration gradient in the sample. Impurities inside the sample may or may not be the cause of the fluctuations of the critical temperature.

The co-existence curve was determined in a large number of series, showing the critical temperature to be constant for each series. Because of the fluctuation of the critical temperature between the individual series, the critical temperature for each series had to be extrapolated from the determined concentrations of the two phases formed; in the case of the succinonitrile-ethanol system, at temperatures below the critical, and in the case of the

Fig. 2. Diffusion constant as a function of the temperature. For the critical system succinonitrile-ethanol the curve is obtained through two different series of experiments. The point marked ? on the critical system triethylamine-water may be due to the difficulties regarding the determination of the critical temperature — see text.

triethylamine-water system, at temperatures above the critical.

The critical temperature, *i.e.*, the temperature at which the first sign of the development of the two phases appeared, was not observed with the requisite accuracy in most of the series. This left an element of doubt regarding the form of the co-existence curve at temperatures closer to the critical, as mentioned above.

The measurement of diffusion constants for the triethylamine-water mixture was carried out in the same way as already reported,⁸ for the mixture of succinonitrile-ethanol at different temperatures from the critical. In Fig. 2 the diffusion constants found are plotted against temperature differences from the critical for both the systems examined. The two figures may permit the assumption that some correlation exists between the dD/dT

of Fig. 2 and the $d\Delta c/d\sqrt[3]{\Delta T}$ of Fig. 1, the latter being the cube root of the constant of formula (1). It must, however, be remembered that the diffusion constants determined concern critical composition, whereas the two phases formed are naturally of compositions different from the critical.

The abscissa scale of Fig. 1 can be determined with greater accuracy than has been done here. However, the primary

aim of the investigations was to study the validity of the "cubic rule"^{1,2} close to the critical point. The determination of the co-existence curves themselves, and the constant of formula (1) for the two mixtures was regarded as being of secondary importance.

The investigation of the triethylamine-water system was complicated by the fact that, at temperatures above, but closer to the critical, the two phases formed did not show the homogeneous composition which is to be expected. Furthermore, at temperatures below the critical, fog was occasionally seen to form inside the prism. After about an hour the fog would disappear, and the section of the concentration curve previously veiled by fog was then seen to have attained an arched form different from the arched concentration curves obtained by diffusion.

The author wishes to thank Professor O. Hassel of the University of Oslo for the interest he has shown in this work.

1. Guggenheim, E. A. *J. Chem. Phys.* **13** (1945) 253.
2. Zimm, B. H. *J. Chem. Phys.* **20** (1952) 538.
3. Lorentzen, H. L. and Bergsnov Hansen, B. *Acta Chem. Scand.* **11** (1957) 893.

Received December 16, 1957.

Reaction Kinetic Determination of the Hydroxyl-Alkoxide Equilibrium in Anion Exchangers

EERO SJÖSTRÖM and LALLI NYKÄNEN

Lääketehtas Orion Oy, Pharmaceutical Manufacturers, Helsinki, Finland

As a sequel to a previous work on the reaction mechanism between estradiol benzoate and an anion exchange resin¹, a kinetic method was developed which is reported here. This method makes it possible to follow the rate of hydrolysis (or alcoholysis) at very low ester concentrations. The method has been used for the investigation of the hydrolysis of methyl benzoate in water and in water-methanol mixtures of low methanol content. The experiments were carried out at $25 \pm 0.02^\circ\text{C}$, by a batch-wise technique with constant stirring. At suitable time

Fig. 1. Weight ratio of methanol and water in the resin phase versus $1/k_{\text{obs}}$ at the concentration range of 0 to 5 % methanol in the outside solution.

intervals samples were taken from the reaction mixture by means of a constant pressure apparatus. The amount of unreacted ester was estimated spectrophotometrically in the ultraviolet region.

According to Samelson and Hammett², the alkaline hydrolysis by anion exchangers in ethanol-water mixtures is a second order reaction. The present results show that when the hydroxyl ion concentration (the resin) is high compared with the ester, the rate constant can be calculated from the first order equation.

On adding methanol to the resin-water mixture the hydroxyl ions are in part converted into alkoxide ions owing to the reaction:

and the hydroxyl ion concentration in the resin phase is decreased. If this reaction is the dominant factor in reducing the hydrolysis rate, it should be possible in theory to calculate the state of this equilibrium from the observed kinetic data. In fact, it was found that at low methanol