

## Occurrence in Plaice Myogen of a Low Molecular Weight Protein of Abnormal Amino Acid Composition

J. W. JEBSEN\* and G. HAMOIR\*\*

Laboratory of General Biology, Institute Van Beneden, University of Liège, Belgium

The ultracentrifugation of the myogens of frog<sup>1</sup>, carp<sup>2</sup> and plaice<sup>3</sup> has shown the presence in these mixtures of a fraction amounting to 40–55 % in the carp and to approximately 25 % in the plaice sedimenting with a rate of about 1.5 Svedberg units which does not occur in rabbit myogen. A component of this fraction precipitating at neutral pH between 90 and 100 % saturation of ammonium sulphate has been crystallized from carp muscle extracts by Henrotte<sup>4</sup> and further investigated by Hamoir<sup>5</sup>. Its ultraviolet absorption appears to be exclusively due to phenylalanine the amount of which has been evaluated spectrophotometrically to about 15 %. We show here that a similar protein exists in plaice myogen too.

This mixture has been fractionated with neutral ammonium sulphate. The fraction precipitating between 90 and 100 % saturation has been examined by electrophoresis in presence of phosphate buffer of  $\mu = 0.05$  and pH = 7.5. A gradient representing about 30 % of the protein content of the fraction and 2 to 3 % of the myogen separates quickly from the slower ones. The corresponding component has been isolated by preparative electrophoresis and examined by ultracentrifugation and spectrophotometry. It sediments as a single peak with a corrected rate of sedimentation of 1.4–1.5 *S*. Its ultraviolet spectrum at pH = 7.5 and 13 is reproduced in Fig. 1. At neutral pH three maxima are observed at 253, 259 and 265  $m\mu$  which correspond closely to those of phenylalanine<sup>6,7</sup>. If the absorption at 259  $m\mu$  is exclusively ascribed to phenylalanine and if use is made of Fromageot and Schneck's<sup>8</sup> data for the absorption of this amino acid, its content can be evaluated to 16 %. A small absorp-




Fig. 1. Ultraviolet spectra at neutral pH (lower curve) and in presence of 0.1 N NaOH (upper curve) of a low molecular weight protein of plaice myogen.

tion of another origin is, however, apparent in both curves at 275–290  $m\mu$ . As it varies from one preparation to the other, it is probably due to an impurity containing tyrosine or cystine. The protein isolated from plaice myogen differs in net charge from the crystalline component found by Henrotte<sup>4</sup>. Its electrophoretic mobility at  $\mu = 0.05$  and pH = 7.5 on the descending side is of  $-6.5 \times 10^{-5}$   $\text{cm}^2 \text{ volt}^{-1} \text{ sec}^{-1}$  while that of the carp protein at  $p = 0.075$  (0.05 in phosphate and 0.025 in NaCl) and pH = 7.3 is of  $-3.20 \times 10^{-5}$   $\text{cm}^2 \text{ volt}^{-1} \text{ sec}^{-1}$ . The content of basic and acid amino acids thus probably varies according to the origin.

An abnormal type of proteins characterized by a low rate of sedimentation, a very high salting-out range, a high content in phenylalanine and the absence of tyrosine and tryptophane appears therefore to occur generally in fish muscle.

\* Norwegian Fisheries Research Institute, Bergen, Norway.

\*\* Associé du F.N.R.S.