On the Crystal Structure of β -ZnOHCl

H.-E. FORSBERG and W. NOWACKI*

Department of Inorganic Chemistry, The Royal Institute of Technology, Stockholm 70, Sweden; Institut für anorganische, analytische und physikalische Chemie, Institut für Mineralogie und Petrographie, Universität, Berne, Switzerland

Recently single crystals of β -ZnOHCl have been prepared from ZnO and ZnCl₂ by Feitknecht and Oswald¹, who kindly supplied us with specimens suitable for X-ray investigations. From rotation and Weissenberg photographs taken around the *a*- and *b*-axes with CuK radiation, it was concluded that the crystals are orthorhombic with the following cell dimensions.

$$a \approx 5.86 \text{ \AA} \quad b \approx 6.58 \text{ \AA} \quad c \approx 11.33 \text{ \AA} \quad V \approx 436.9 \text{ \AA}^3$$

The unit cell contains eight formula units.

The extinctions *h*0*l* for *h* odd, *hk*0 for *k* odd and 0*kl* for *l* odd are in agreement with the centrosymmetric space group *Pcab* (No. 61 in the *International Tables for X-ray Crystallography*²) which also has been confirmed by the structure determination.

The following preliminary parameters of the atoms were determined mainly from Patterson and Fourier projections of *h*0*l* and 0*kl*:

8 Zn in 8(c) with $x = 0.250$ $y = 0.131$ $z = 0$

8 O in 8(c) with $x = 0.594$ $y = 0.126$ $z = 0.071$

8 Cl in 8(c) with $x = 0.583$ $y = 0.107$ $z = 0.357$

8(c) $\pm (x, y, z)$, $\pm (\frac{1}{2} - x, \frac{1}{2} + y, \bar{z})$,
 $\pm (\frac{1}{2} + x, \bar{y}, \frac{1}{2} - z)$, $\pm (x, \frac{1}{2} - y, \frac{1}{2} + z)$.

Figs. 1 and 2 show the structure of β -ZnOHCl. The zinc atoms are surrounded by three oxygen and three chlorine atoms, so that a coordination Zn-6X in the shape of an octahedron is obtained. The structure can also be described as pseudohexagonal zinc layers separated by an ordered layer of oxygen and chlorine atoms.

* Communication No. 110 of Abteilung für Kristallographie und Strukturlehre, Mineralogisches Institut Universität Bern.

Fig. 1. The structure of β -ZnOHCl, viewed along [100]. (The range for the *x* parameter is $-0.406 \leq x \leq 0.594$.)

Fig. 2. The figure shows the pseudohexagonal zinc layer at $z = \frac{1}{2}$.

The following ranges of interatomic distances were found.

Zn—Zn	3.29—3.40 Å
Zn—O	2.01—2.17 Å
Zn—Cl	2.46—2.56 Å
O—O	2.56—3.36 Å
O—Cl	3.08—3.51 Å
Cl—Cl	3.48—4.08 Å

The complete structure determination will be published later.

1. Feitknecht, W. and Oswald, H. R. *Private communication*.
2. *International Tables for X-ray Crystallography*, Kynoch Press, Birmingham 1952, Vol. I, p. 150.

Received June 12, 1959.

Unit Cell and Space Group of VS_4

BJØRN PEDERSEN

*Kjemisk institutt, Universitetet i Oslo,
Blindern, Norway*

In an X-ray and magnetic study of the system V/S by Klemm and Hoschek¹ a polysulphide of vanadium was discovered. By means of vapour pressure measurements the composition of this compound was decisively determined as VS_4 by Biltz and Köcher². A polysulphide of vanadium is also found in nature as patronite first described by Hewett³. On the basis of analytical data given by Hillebrand⁴, Biltz and Köcher² concluded that patronite is VS_4 . A sample of this mineral was kindly placed at our disposal by the Curator of Harvard Museum, Dr. C. Frondel. Powder photographs showed that the major phase present was the same as synthetic VS_4 . — No crystallographic data on VS_4 seems to have been reported.

In a recent investigation of the vanadium sulphides⁵, single crystals of VS_4 were obtained by heating a mixture of $VS_{1.5}$ and sulphur, corresponding to the formula VS_4 , in an evacuated and sealed silica tube for four months at 400°C and slowly cooled to 100°C over another four months. The reaction was not complete, however, as some sulphur was still present in the tube. The crystals had grown as clustres of long, black needles with metallic lustre. Most of them were unsuitable for X-ray work, but after some search a crystal with very good properties was found. Its dimensions were: $0.33 \times 0.06 \times 0.07$ mm³.

Oscillation and Weissenberg photographs taken with CuK -radiation ($\lambda_{Cu} = 1.5405$ Å)

around the needle axis (the c -axis) showed the crystal to be monoclinic. This was confirmed by precession photographs taken with MoK -radiation of the $h0l$ -plane. From Guinier powder photographs the cell dimensions were found to be:

$$a = 12.67 \text{ Å} \quad b = 10.41 \text{ Å} \quad c = 12.11 \text{ Å} \\ \beta = 148.37^\circ$$

Missing reflexions were: hkl when $h+k = 2n+1$, $h0l$ when $l = 2n+1$ and $(h = 2n+1)$ and $0k0$ when $(k = 2n+1)$. This is characteristic of the space groups No. 9, Cc and No. 15, $C2/c$. The density was determined by flotation to be 2.80 g/cm³, which indicates the presence of eight VS_4 groups in the unit cell ($d_{calc} = 2.835$ g/cm³).

As the basis for a structure determination, the intensities of the $hk0$ - and $h0l$ -reflexions were estimated visually. The intensities were corrected for Lorentz and polarisation effects, but not for absorption and extinction.

This intensity material was used to decide between the two characteristic space groups by means of the Wilson ratio for detection of centers of symmetry, $\rho = \langle |F_o|^2 \rangle^2 / \langle |F_c|^2 \rangle^2$. For the 48 $hk0$ - and 38 $h0l$ -reflexions ρ was calculated as 0.77 and 0.71, respectively. The predicted value for ρ is 0.635 for centrosymmetric crystals and 0.785 for noncentrosymmetric crystals. Thus the Wilson ratio indicates that the crystal is noncentrosymmetric and the space group accordingly No. 9, Cc .

Some crystals were kindly tested for piezoelectricity by Dr. W. G. Perdok of the University of Groningen, but with negative results. This might perhaps be due to the smallness of the crystals (< 0.06 mm) in the direction of the polar axis.

1. Klemm, W. and Hoschek, E. *Z. anorg. u. allgem. Chem.* **226** (1936) 359.
2. Biltz, W. and Köcher, A. *Z. anorg. u. allgem. Chem.* **242** (1939) 324.
3. Hewett, F. *Eng. Mining J.* **1906** 385.
4. Hillebrand, W. F. *J. Am. Chem. Soc.* **29** (1912) 1019.
5. Pedersen, B. *Thesis*, University of Oslo, 1958.
6. Wilson, A. J. C. *Acta Cryst.* **2** (1949) 318.

Received June 5, 1959.