5. Westöb, G. *Acta Chem. Scand.* **13** (1959) 604.
6. Pepinsky, R., Robertson, J. M. and Speakman, J. C. *Computing Methods and the Phase Problem in X-ray Crystal Analysis*, Pergamon Press, Oxford 1961; Rollett, J.S., p. 87; Sime, J. G. p. 301.

Received March 5, 1962.

On the Structure of CrOCl

HANS-ERIK FORSBERG

Department of Inorganic Chemistry, The Royal Institute of Technology, Stockholm 70, Sweden

Some years ago single crystals of CrOCl were prepared from Cr₂O₃ and CrCl₃ by Professor H. Schäfer¹, who was kind enough to supply me with specimens suitable for X-ray investigations. From rotation and Weissenberg photographs taken around the *b*-axis with CuK radiation, it was concluded that the crystals are orthorhombic with the following cell dimensions:

$$a \approx 3.88 \text{ \AA}, b \approx 3.20 \text{ \AA}, c \approx 7.72 \text{ \AA}, \\ V \approx 95.9 \text{ \AA}^3.$$

From the observed density of 3.47 g/ml it was calculated that there should be two formula units of CrOCl in the unit cell.

The extinctions, $hk0$ for $h + k = \text{odd}$, are in agreement with the centrosymmetric space group $Pm\bar{m}n$ and the non-centrosymmetric space group $Pmn2_1$. (Nos. 59 and 31 in the *International Tables for X-ray Crystallography*²). Since a plausible structure could be assigned assuming $Pm\bar{m}n$, the space group of higher symmetry, the lower space group was not considered.

The following preliminary parameters of the atoms were determined from Patterson and Fourier projections on the xz plane and geometrical considerations:

- 2 Cr in $2(a)$: $\pm (1/4, 1/4, z)$ with $z = 0.109$
 2 Cl in $2(b)$: $\pm (1/4, 3/4, z)$ with $z = 0.327$
 2 O in $2(b)$: $\pm (1/4, 3/4, z)$ with $z = 0.96$

Fig. 1 shows the structure of CrOCl. The chromium and oxygen atoms form Cr—O—Cr layers which are separated by double halogen sheets.

- ○ ⊖ are Cr, O and Cl at $y = 1/4$.
 ● ⊖ ⊖ are Cr, O and Cl at $y = 3/4$.

Fig. 1. The structure of CrOCl, viewed along $[010]$.

The atomic arrangement and the space group of CrOCl is the same as in FeOCl (Goldsztaub³), InOCl and InOBr (Forsberg⁴), and TiOCl (Schäfer⁵). Thus, as postulated by Schäfer, chromium oxide chloride is isomorphous with these substances.

The following ranges of interatomic distances were found:

Cr—Cr	3.04 Å
Cr—O	1.97–2.01 Å
Cr—Cl	2.32 Å
O—O	2.51 Å
O—Cl	2.83–3.35 Å
Cl—Cl	3.20–3.88 Å

A complete structure determination will be published later.

The research reported in this document has been sponsored by Air Force Office of Scientific Research, OAR, through the European Office, Aerospace Research, United States Air Force, under contract No. AF 61(052)-162.

1. Schäfer, H. and Wartenpfehl, F. *Z. anorg. u. allgem. Chem.* **308** (1961) 282.
2. *International Tables for X-ray Crystallography*, Kynoch Press, Birmingham 1952, Vol I, p. 148 and 117.
3. Goldsztaub, M. S. *Compt. rend.* **198** (1934) 667.
4. Forsberg, H.-E. *Acta Chem. Scand.* **10** (1956) 1286.
5. Schäfer, H., Wartenpfehl, F. and Weise, E. *Z. anorg. u. allgem. Chem.* **295** (1958) 268.

Received March 8, 1962.