Replacement of the purified Baranowski enzyme in the above system by cell-sap, obtained from the preparation of the rat skeletal muscle mitochondria, also resulted in an enhanced DPNH oxidation, which again was sensitive to antimycin A and insensitive to amytal and rotenone. 1.1 mg cell-sap protein per mg mitochondrial protein gave rise to a glycerol-1-phosphate cycle activity of about 50 %, and 3.7 mg cell-sap protein to an activity of almost 100 %, of that obtained with an excess of purified Baranowski enzyme.

These results demonstrate the operation of the glycerol-1-phosphate cycle in a reconstructed subcellular system from rat skeletal muscle. The contribution of this cycle to the oxidation of extramitochondrial DPNH by the mitochondrial respiratory chain in the intact muscle may be ascertained in the future by investigating the effects of antimycin A, amytal and rotenone on the respiration of whole muscle under appropriate experimental conditions.

This work has been supported by a grant from the *Swedish Medical Research Council*.

1. Hedman, R., Suranyi, E. M., Luft, R. and Ernster, L. *Biochem. Biophys. Res. Commun.* **8** (1962) 314.
2. Bücher, T. and Klingenberg, M. *Angew. Chem.* **70** (1958) 552.
3. Estabrook, R. W. and Sacktor, B. *J. Biol. Chem.* **233** (1958) 1014.
4. Zebe, E., Delbrück, A. and Bücher, T. *Biochem. Z.* **331** (1959) 254.
5. Sacktor, B. and Dick, A. *J. Biol. Chem.* **237** (1962) 3259.
6. Borst, P. *Biochim. Biophys. Acta* **57** (1962) 270.

Steroid Effects on Antimycin Inhibited NADH and Succinate Oxidation

P. K. JENSEN

Division of Endocrinology, Institute for Experimental Medicine, University of Copenhagen, Copenhagen, Denmark

In experiments where spectral changes in the respiratory chain components of heart sarco-
somal fragments were recorded during oxida-

tion of NADH, it was observed that the time required for the suspension to expend dissolved oxygen in the presence of antimycin was longer than when deoxycorticosterone was added together with antimycin. Deoxycorticosterone and other steroids are potent inhibitors of NADH oxidation^{1,2} and although Yielding *et al.*³ found that the steroids apparently lost their inhibitory effect in the presence of antimycin, a partial reversal of antimycin inhibition as indicated here was unexpected.

A change in the rate of oxygen consumption could be demonstrated directly in experiments where oxygen concentration was recorded by means of a Clark oxygen electrode. The combined addition of antimycin and deoxycorticosterone led to a rate of oxygen consumption lower than that obtained with steroid alone, but higher than that with antimycin added alone. It appeared as if the steroid had prevented antimycin from exerting its full inhibitory effect. Preincubation of ETP with antimycin prior to the start of the reaction and addition of steroid did not abolish this effect of the steroid.

Neither steroids nor amytal normally inhibit succinate oxidation, and this reaction should therefore provide a more simple system for investigation of steroid and amytal effects on antimycin inhibition. When NADH was replaced with succinate and oxygen concentration recorded, the results showed, however, that steroid had an effect on antimycin inhibited succinate oxidation opposite to what it had on antimycin inhibited NADH oxidation. Amytal in 20 times higher concentrations had no effect at all. Again preincubation of ETP with antimycin did not change this picture. The results with succinate were, however, less reproducible than with NADH. Ageing both at -15° and at 0°C appeared to influence the response to steroid, and also the initial succinate concentration in the reaction medium played a role.

These observations show that steroid and amytal can react with the respiratory chain in a more complex manner than previously assumed, and that steroid and amytal need not necessarily have identical effects.

1. Jensen, P. K. *Nature* **184** (1959) 451.
2. Yielding, K. L. and Tomkins, G. M. *Proc. Natl. Acad. Sci. U.S.A.* **45** (1959) 1730.
3. Yielding, K. L., Tomkins, G. M., Munday, J. S. and Cowley, I. J. *J. Biol. Chem.* **235** (1960) 3413.