

4. Woldbye, F. *Optical Rotatory Dispersion of Transition Metal Complexes*. European Research Office, U. S. Army, Frankfurt a.M. 1959.
5. Fronåus, S. *Komplexsystem hos Koppar*, *Diss.* Lund 1948.
6. Cannan, R. K. and Kibrick, A. *J. Am. Chem. Soc.* **60** (1938) 2314.
7. Bjerrum, J. and Larsen, E. in *Essays in Coordination Chemistry*. Birkhauser Verlag, Basel 1964.
8. Theilacker, W. *Z. anorg. Chem.* **234** (1937) 161.
9. Mathieu, J.-P. *Ann. Phys. Paris* 11. serie **19** (1944) 335.
10. Corey, E. J. and Bailar, J. C. *J. Am. Chem. Soc.* **81** (1959) 2620.
11. Marckwald, W. *Ber.* **37** (1904) 1038.
12. Thomé, L. G. *Ber.* **36** (1903) 582.
13. Lindhard, M. and Weigel, M. *Z. anorg. Chem.* **260** (1949) 65.
14. Lindhard, M. and Rau, B. *Z. anorg. Chem.* **271** (1953) 121.

Received April 22, 1964.

## Effect of Long Heat-Denaturation of Collagen on Its Electrophoretic Pattern

T. HOLLMÉN and E. KULONEN

Department of Medical Chemistry, University of Turku, Turku 3, Finland

For the denaturation of soluble collagen, Piez *et al.*<sup>1</sup> suggested heating at pH 4.8 for 15 min at +40°C. The detailed investigation by Engel<sup>2</sup> showed that the denaturation proceeds in two stages: the helical structures are lost very rapidly but the separation of the chains takes more time, and he suggested heating for 90 min for complete denaturation. In a later paper Engel and Beier<sup>3</sup> demonstrated that the  $\alpha$ - and  $\beta$ -components are not stable, but already after 9 h at +40°C in pH 3.7 citrate buffer 10–20 % of the  $\beta$ -component was lost from the sedimentation pattern, and also degradation products of small molecular weight were obtained from both  $\alpha$ - and  $\beta$ -fractions. These findings prompted us to check the electrophoretic pattern of heat-denatured collagen after various intervals.

The collagen preparation was obtained from rat tail tendon fibres by extraction for 24 h in the cold with 20 volumes of 3 % (v/v) acetic acid. (The material had been first extracted exhaustively with 0.4 M NaCl, buffered to pH 7.3 with 1/15 M phosphate, and the soluble fraction discarded.) The collagen was precipitated from the acetic acid extract by dialysis against pH 7.3, M/15 phosphate buffer. The precipitate was lyophilized, dissolved in 3 % acetic acid and dialyzed against the pH 4.5 acetate buffer, ionic strength  $\mu = 0.022$ , which was also used in the preparation of the gel for the electrophoretic run.<sup>4</sup> For the prevention of microbial growth, the buffer was saturated with octanol. No turbidities were observed. After a preliminary denaturation at +40°C for 15 min, the sample was divided in separate lots, which were kept at +4°C, until they




Fig. 1. Effect of long heat-denaturation of soluble collagen on the starch gel electrophoretic pattern. The duration of the denaturation at +40°C is indicated. The sample size was 400  $\mu$ g, but in some runs larger samples were applied to accentuate the fast-moving fraction F (the pattern to the extreme right).

were immersed in a +40°C water bath at appropriate intervals to get the indicated heating times at the beginning of the simultaneous electrophoretic runs on the same gel sheet (Fig. 1, except the run marked 5 d-B.).

The pattern does not change essentially in a week, but three observations were made: (1) The slow-migrating fractions were gradually lost during the heating and they therefore represent larger complexes, which are broken down. (2) The  $\alpha_1$ -fraction becomes rather broad. The separation to subcomponents is not clear, but on the basis of unpublished work by V. Nántö from our laboratory we believe that this fraction is not homogeneous. (3) After about 3 days there appears a new distinct band designated F, which migrates faster than the  $\alpha_2$ -component. In Fig. 1 the

run 5d—B is included to demonstrate this fraction more clearly. Also in the preparative modification of the starch gelelectrophoresis (unpublished work) there appears a fast-moving fraction of gelatin. The migration rate of the new band corresponds to that of the commercial gelatin obtained from limed precursors.<sup>4</sup> The possibility of an artefact from microbial growth does not seem likely to us, mainly because of the regular appearance and distinct form of the band, which has been observed also in the fractionation of other samples of denatured collagen.

This work forms a part of a program supported by institutional grants from *U. S. Department of Agriculture*, Foreign Research and Technical Programs Division, and from *Sigröd Jusélius Foundation*.

1. Piez, K. A., Weiss, E. and Lewis, M. S. *J. Biol. Chem.* **235** (1960) 1987.
2. Engel, J. *Arch. Biochem. Biophys.* **97** (1962) 150.
3. Engel, J. and Beier, G. *Z. physiol. Chem.* **334** (1963) 201.
4. Nántö, V., Maatela, J. and Kulonen, E. *Acta Chem. Scand.* **17** (1963) 1604.

Received May 14, 1964.

## Torsional Coordinate in the Book of Wilson, Decius and Cross

S. J. CYVIN and J. BRUNVOLL

*Institute of Theoretical Chemistry, Technical University of Norway, Trondheim, Norway*

During a work on a computer programme for the general B matrix in the theory of molecular vibrations,<sup>1</sup> the torsional coordinates showed a wrong behaviour. The origin of this mistake was unexpected, as it was found to be an inexactness in the book

of Wilson, Decius and Cross,<sup>2</sup> from which we had taken the expressions on p. 61. The error should be corrected since this excellent book represents one of the standard references of all investigators in the field of molecular spectroscopy.

The expression (22) on p. 61 (Ref.<sup>2</sup>) is not general. In particular it is observed that it does not contain the equilibrium dihedral angle  $\tau$ . The expression was found to apply only to the special planar case of  $\tau = \pi$  (but *not*  $\tau = 0$ ). The torsional coordinates of the benzene ring ( $\tau = 0$ ) given on p. 256 showed to be correct, but they do *not* follow from (21) — (24), as stated in the text. On the other hand, they were found to be completely consistent with the correct general expressions given by Decius in an earlier publication.<sup>3</sup> For convenience these expressions are given below, as translated to the notation of Wilson, Decius and Cross.<sup>2</sup>

$$\begin{aligned}
 s_{t1} &= -\frac{\mathbf{e}_{12} \times \mathbf{e}_{23}}{r_{12} \sin^2 \varphi_2} \\
 s_{t2} &= \left[ \frac{r_{23} - r_{12} \cos \varphi_2}{r_{23} r_{12} \sin \varphi_2} - \frac{\cos \tau \cos \varphi_3}{r_{23} \sin \varphi_3} \right] \frac{\mathbf{e}_{12} \times \mathbf{e}_{23}}{\sin \varphi_2} \\
 &\quad - \frac{\sin \tau \cos \varphi_3}{r_{23} \sin \varphi_3} \mathbf{e}_{23} \times \left( \frac{\mathbf{e}_{12} \times \mathbf{e}_{23}}{\sin \varphi_2} \right) \\
 s_{t3} &= [(14)(23)]s_{t2} \\
 s_{t4} &= [(14)(23)]s_{t1}
 \end{aligned}$$

In this connection we also wish to report some rules for the direction of movement for the atoms during the torsional displacement. When it is adhered to the usual sign convention and consecutive numbering of atoms,<sup>2</sup> one has: For positive  $\Delta\tau$  atom 1 will move in the direction of the vector  $\mathbf{e}_{21} \times \mathbf{e}_{23}$ , and atom 4 in the direction of  $\mathbf{e}_{34} \times \mathbf{e}_{32}$ .

1. Cyvin, S. J. and Brunvoll, J. *Acta Polytechn. Scand. In press.*
2. Wilson, E. B., Jr., Decius, J. C. and Cross, P. C. *Molecular Vibrations*, McGraw-Hill Book Co., New York 1955.
3. Decius, J. C. *J. Chem. Phys.* **16** (1948) 1025.

Received April 14, 1964.